

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO. (If known see 37 C.F.R. 1.5)

09/674817

INTERNATIONAL APPLICATION NO.

PCT/EP99/03141

INTERNATIONAL FILING DATE

7 May 1999

(EARLIEST) PRIORITY DATE CLAIMED

8 May 1998

TITLE OF INVENTION

NUCLEIC ACID MOLECULES WHICH CODE FOR ENZYMES DERIVED
FROM WHEAT AND WHICH ARE INVOLVED IN THE SYNTHESIS OF
STARCH

APPLICANTS FOR DO/EO/US

Horst LÖRZ, Stephanie LÜTTICKE, Gernot ABEL and
Ulrich GENSCHER

Applicants herewith submit to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)) including the International Search Report and the amended pages:
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)), including ___ sheets of formal drawings and a copy of the International Search Report.
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventors (35 U.S.C. 371(c)(4)).
10. ☐ The annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern other document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
PCT/ISA/210, PCT/IB/306, 308
PCT/RO/101, 18 pages of Sequence Listing


EXPRESS MAIL

Mailing Label Number: EL585029828US


Date of Deposit: November 6, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service

"Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents and Trademarks, Box PCT Washington, DC 20231.



(Typed or printed name of person mailing paper or fee)


(Signature of person mailing paper or fee)

17. ☒ The following fees are submitted:

(CALCULATIONS /PTO USE ONLY)

Basic National Fee (37 CFR 1.492(a)(1)-(5):

Neither international preliminary examination fee (37 CFR 1.482) nor
international search fee (37 CFR 1.445(a)(2)) paid to USPTO and
International Search Report not prepared by the EPO or JPO... \$1000.00

International preliminary examination fee (37 C.F.R. 1.482)
not paid to USPTO but International Search Report prepared
by the EPO or JPO.....\$ 860.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
Not paid to USPTO but international search fee (37 CFR 1.445(a)(2))
paid to USPTO (37 CFR 1.445(a)(2)).....\$ 710.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
But all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$ 690.00

International preliminary examination fee paid to USPTO (37 CFR 1.482)
and all claims satisfied provisions of PCT Article 33(1)-(4).....\$ 100.00

ENTER APPROPRIATE BASIC FEE AMOUNT = (\$ 860.00)

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(e)). (\$ /

24 Claims	/Number Filed	/	Number Extra	/Rate	(
Total Claims	/ 24 - 20 =	/	4	X \$18.00	\$ 72.00

Independent Claims	/ 1 - 3 =	/	0	/X \$80.00	(\$ 0.00
--------------------	-----------	---	---	------------	----------

Multiple dependent claim(s) (if applicable)	/+ \$260.00	(\$
---	-------------	-----

TOTAL OF ABOVE CALCULATIONS = (\$ 932.00

Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity (statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28) (/

SUBTOTAL = (\$ 932.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30 months from the earliest claimed priority date (37 CFR 1.492(f)).+ (\$ /

TOTAL NATIONAL FEE = (\$ 932.00

Fee for recording the enclosed assignments (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) \$40.00 per property + (\$ 40.00/ (\$ /

TOTAL FEES ENCLOSED = (\$ 972.00

(Amount to be: /
(refunded /\$
(charged /\$

- a. ☒ Our checks in the amount of \$972.00 to cover the above fees is enclosed.
b. ☐ Please charge my Deposit Account No. 50-0320 in the amount of \$ to cover the above fees.
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 50-0320. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

**WILLIAM F. LAWRENCE
FROMMER LAWRENCE & HAUG LLP
745 FIFTH AVENUE
NEW YORK, NEW YORK 10151**

SIGNATURE

**WILLIAM F. LAWRENCE
NAME**

Dated: November 6, 2000

28,029
REGISTRATION NUMBER

00907 47842950

09/674817

529 Rec'd PCT/PTO 06 NOV 2000
514413-3849

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) : LÖRZ et al.
U.S. Serial No. : Filed Concurrently Herewith
Int'l Appln. No. : PCT/EP99/03141
Int'l Filing Date : 07 May 1999
Earliest Priority Date : 08 May 1998
For : **NUCLEIC ACID MOLECULES WHICH CODE FOR
ENZYMES DERIVED FROM WHEAT AND WHICH
ARE INVOLVED IN THE SYNTHESIS OF STARCH**


745 Fifth Avenue
New York, NY 10151


EXPRESS MAIL

Mailing Label Number: EL585029828US

Date of Deposit: November 6, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" Service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Box PCT, Washington, D.C. 20231.


(Typed or printed name of person
mailing paper or fee)


(Signature of person mailing paper or fee)

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Box PCT
Washington, D.C. 20231

Dear Sir:

Preliminary to the examination of this U.S. national phase application, please enter the following amendments:

09674817 4106000

IN THE CLAIMS:

Please cancel Claims 1-26 and replace them with Claims 27-49:

27. A nucleic acid molecule encoding a protein with the function of a wheat isoamylase, selected from the group consisting of
- (a) a nucleic acid molecule encoding a protein comprising the amino acid sequence shown under Seq ID No. 3,
 - (b) a nucleic acid molecule comprising the nucleotide sequence shown under Seq ID No. 2 or a part thereof or a ribonucleotide sequence corresponding hereto;
 - (c) a nucleic acid molecule which hybridizes with a nucleic acid molecule mentioned under (a) or (b) or is complementary thereto, and
 - (d) a nucleic acid molecule whose nucleotide sequence deviates from the sequence of a nucleic acid molecule mentioned under (a), (b) or (c) owing to the degeneracy of the genetic code,
- the nucleic acid molecule mentioned under (a), (c) and (d) having a homology of over 90% with Seq ID No. 2.
28. A nucleic acid molecule as claimed in claim ₂₇ which is a DNA molecule.
29. A DNA molecule as claimed in claim ₂₈ which is a cDNA molecule.
30. A nucleic acid molecule as claimed in claim ₂₇ containing regulatory elements.
31. A nucleic acid molecule as claimed in claim ₂₇ which is an RNA molecule.
32. A nucleic acid molecule which specifically hybridizes with a nucleic acid molecule as claimed in claim ₂₇ and has a homology of over 90% with Seq ID No. 2.
33. A nucleic acid molecule as claimed in claim ₃₂ which, is an oligonucleotide with a length of at least 15 nucleotides.

34. A vector containing a DNA molecule as claimed in claim 27
35. A vector as claimed in claim 34 wherein said nucleic acid molecule is linked in sense orientation to regulatory elements which ensure transcription and synthesis of a translatable RNA in pro- or eukaryotic cells.
36. A vector as claimed in claim 34 wherein said nucleic acid molecule is linked in sense orientation to regulatory elements which ensure the synthesis of an untranslatable RNA in pro- or eukaryotic cells.
37. A vector as claimed in claim 34 wherein said nucleic acid molecule is linked in antisense orientation to regulatory elements which ensure the synthesis of an untranslatable RNA in pro- or eukaryotic cells.
38. A host cell which is transformed with a nucleic acid molecule as claimed in one or more of claims 27 to 31 or a vector as claimed in one or more of claims 34 to 37 or which is derived from such a cell.
39. A process for the preparation of a protein as claimed in claim 39 wherein a host cell as claimed in claim 38 is cultured under conditions which permit said protein to be synthesized and said protein is isolated from the cultured cells and/or the culture medium.
40. A process for generating a transgenic plant cell, wherein
a) a nucleic acid molecule as claimed in one or more of claims 27 to 31
or
b) a vector as claimed in one or more of claims 34 to 37
is integrated into the genome of a plant cell.
41. A transgenic plant cell which has been transformed with a nucleic acid molecule as claimed in one or more of claims 27 to 31 or one or more vector as claimed in claims 34 to 37 or which is derived from such a cell.
42. A process for generating a transgenic plant cell, wherein

a1) a nucleic acid molecule as claimed in one or more of claims 27 to 31
or

a2) a vector as claimed in one or more of claims³⁴ to ³⁷
is integrated into the genome of a plant cell and

b) an intact plant is regenerated from said plant cell.

43. A plant containing a plant cell as claimed in claim 41.

44. A plant as claimed in claim 43 which is a crop plant.

45. A plant as claimed in claim 44 which is a starch-storing plant.

46. A plant as claimed in claim 45 which is a monocotyledonous plant
or maize

47. A plant as claimed in claim 46 which is a barley, rye or wheat plant.

48. A propagation material of a plant as claimed in one or more of
claims₄₃ to₄₇ .

49. The use of a plant cell as claimed in claim 41, a plant as claimed in
one or more of claims ₄₃ to ₄₇ or of propagation material as claimed
in claim₄₈ for the production of starch. ---


009077" 27842950
09674817 110600

REMARKS

Entry of this amendment and an early examination on the merits are respectfully
solicited.

Respectfully submitted,

FROMMER LAWRENCE & HAUG LLP

By: 
William F. Lawrence
Reg. No. 28,029
(212) 588-0800

009077 2787950

WO 99/58690

PCT/EP99/03141

Nucleic acid molecules encoding wheat enzymes involved in starch synthesis

5 The present invention relates to nucleic acid molecules which encode a wheat enzyme involved in starch synthesis in plants. This enzyme is an isoamylase.

10 The invention furthermore relates to vectors, host cells, plant cells and plants comprising the nucleic acid molecules according to the invention.

Furthermore, there are described methods for the generation of transgenic plants which, owing to the introduction of nucleic acid molecules according to the invention, synthesize starch with altered characteristics.

15 In view of the increasing importance attributed lately to plant constituents as renewable raw materials, one of the objects of biotechnology research addresses the adaptation of these plant raw materials to the needs of the processing industries. Moreover, to allow renewable raw materials to be used in as many fields as possible, a wide diversity of materials must be generated.

20 Apart from oils, fats and proteins, polysaccharides constitute the important renewable raw materials from plants. Apart from cellulose, starch – which is one of the most important storage substances in higher plants – takes a central position amongst the polysaccharides. In this context, wheat is one of the most important crop plants since it provides approximately 20% of the total starch production in the European Community.

30 The polysaccharide starch is a polymer of chemically uniform units, the glucose molecules. However, it is a highly complex mixture of different molecule types which differ with regard to their degree of polymerization, the occurrence of branching of the glucose chains and their chain lengths, which, in addition, may be derivatized, for example phosphorylated. Starch therefore does not constitute a uniform raw material. In particular, a distinction is made between amylose starch, an essentially unbranched polymer of alpha-1,4-glycosidically linked glucose molecules, and amylopectin starch, which, in turn, constitutes a complex mixture of glucose chains with various branchings. The branchings occur by the occurrence of

09674817-110600

additional alpha-1,6-glycosidic linkages. In wheat, amylose starch makes up approximately 11 to 37% of the starch synthesized.

To allow suitable starches to be used in the widest possible manner for the widest possible range of industrial needs, it is desirable to provide plants which are capable of synthesizing modified starches which are particularly well suited to various purposes. One possibility of providing such plants is to employ plant-breeding measures. However, since wheat is polyploid in character (tetra- and hexaploid), the exertion of influence by plant breeding proves to be very difficult. A "waxy" (amylose-free) wheat was generated only recently by crossing naturally occurring mutants (Nakamura et al., Mol. Gen. Genet. 248 (1995), 253-259).

An alternative to plant-breeding methods is the specific modification of starch-producing plants by recombinant methods. However, prerequisites are the identification and characterization of the enzymes which are involved in starch synthesis and/or starch modification and the isolation of the nucleic acid molecules encoding these enzymes.

The biochemical pathways which lead to the synthesis of starch are essentially known. Starch synthesis in plant cells takes place in the plastids. In photosynthetically active tissue, these plastids are the chloroplasts and in photosynthetically inactive, starch-storing tissue are amyloplasts.

A further specific alteration of the degree of branching of starch synthesized in plants with the aid of recombinant methods still requires identification of DNA sequences, which encode enzymes involved in starch metabolism, in particular in the introduction or degradation of branching within the starch molecules.

Besides the so-called Q enzymes, which introduce branchings into starch molecules, enzymes occur in plants which are capable of breaking down branchings. These enzymes are called debranching enzymes and, according to their substrate specificity, they are divided into three groups:

- (a) The pullulanases, which, in addition to pullulan, also utilize amylopectin as substrate, are found in microorganisms, for example *Klebsiella* and in plants. In plants, these enzymes are also termed R enzymes.

- (b) The isoamylases, which do not utilize pullulan, but indeed glycogen and amylopectin as substrate, are also found in microorganisms and plants. For example, isoamylases have been described in maize (Manners & Carbohydr. Res. 9 (1969), 107) and potato (Ishizaki et al., Agric. Biol. Chem. 47 (1983), 771-779).
- (c) The amylo-1,6-glucosidases are described in mammals and yeasts and utilize grenzdextrins as substrate.

In sugar beet, Li et al. (Plant Physiol. 98 (1992), 1277-1284) were only able to find one debranching enzyme of the pullulanase type, in addition to five endoamylases and two exoamylases. This enzyme, which has a size of approx. 100 kD and a pH optimum of 5.5, is localized in the chloroplasts. In spinach, too, a debranching enzyme was described which utilizes pullulan as substrate. The activity both of the spinach debranching enzyme and of the sugar beet debranching enzyme upon reaction with amylopectin as substrate is five times lower in comparison with pullulan as substrate (Ludwig et al., Plant Physiol. 74 (1984), 856-861; Li et al., Plant Physiol. 98 (1992), 1277-1284).

In the agronomically important starch-storing crop plant potato, the activity of a debranching enzyme was studied by Hobson et al. (J. Chem. Soc., (1951), 1451). It was proved successfully that, in contrast to the Q enzyme, this enzyme has no chain-extending activity, but merely hydrolyzes alpha-1,6-glycosidic bonds. However, it has been impossible as yet to characterize the enzyme in greater detail. In the case of potatoes, processes for purifying the debranching enzyme and partial peptide sequences of the purified protein have already been proposed (WO 95/04826). In the case of spinach, the purification of a debranching enzyme and the isolation of suitable cDNA have been described in the meantime (Renz et al., Plant Physiol. 108 (1995), 1342).

In maize, only the existence of one debranching enzyme has been described as yet in the literature. Owing to its substrate specificity, this enzyme is classified as belonging to the group of the isoamylases (see, for example, Hannah et al., Scientia Horticulturae 55 (1993), 177-197 or Garwood (1994) in Starch Chemistry and Technology, Whistler, R.L., BeMiller, J.N., Puschall, E.F. (eds.), Academic Press San Diego, New York, Boston, 25-86). The corresponding mutant is termed "sugary". The gene of the sugary locus has been cloned recently (see James et al., Plant Cell 7

(1995), 417-429). Apart from the sugary locus, no other gene locus which encodes a protein with debranching enzyme activity is as yet known in maize. Also, there have been no indications to date that other debranching enzyme forms occur in maize. If transgenic maize plants are to be generated which no longer have any debranching enzyme activities whatsoever, for example in order to extend the degree of branching of the amylopectin starch, it is necessary to identify all debranching enzymes forms which occur in maize and to isolate the corresponding genes or cDNA sequences.

To provide further possibilities of altering any starch-storing plant, preferably cereals, in particular wheat, so that it synthesizes a modified starch, it is necessary to identify in each case DNA sequences which encode further isoforms of branching enzymes.

The object of the present invention is therefore to provide nucleic acid molecules encoding enzymes involved in starch synthesis, which allow genetically modified plants to be generated which make possible the production of plant starches whose chemical and/or physical characteristics are altered.

This object is achieved by providing the use forms designated in the patent claims.

The present invention therefore relates to a nucleic acid molecule which encodes a protein with the function of a wheat isoamylase, preferably a protein which is essentially defined by the amino acid sequence stated under Seq ID No. 3 or 7. In particular, the invention relates to a nucleic acid molecule comprising the nucleotide sequence stated under Seq ID No. 1, 2 or 6, or a part thereof, preferably a molecule comprising the coding region stated in Seq ID No. 1, 2 or 6, and corresponding ribonucleotide sequences. Very especially preferred is a nucleic acid molecule furthermore comprising regulatory elements which ensure transcription and, if appropriate, translation of said nucleic acid molecules. The subject matter of the invention is furthermore a nucleic acid molecule which hybridizes with one of the nucleic acid molecules according to the invention.

The subject matter of the invention is also a nucleic acid molecule encoding a wheat isoamylase whose sequence deviates from the nucleotide sequences of the above-described molecules owing to the degeneracy of the genetic code.

5

The invention also relates to a nucleic acid molecule with a sequence which is complementary to all or part of one of the abovementioned sequences.

- 10 The term "hybridization" as used in the context of the present invention denotes hybridization under conventional hybridization conditions, preferably under stringent conditions, as they are described, for example, by Sambrook et al., Molecular Cloning, A Laboratory Manual, 2nd Ed. (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

15

"Hybridization" especially preferably takes place under the following conditions:

Hybridization buffer: 2 × SSC; 10 × Denhardt solution (Fikoll 400 + PEG + BSA; ratio 1:1:1); 0.1% SDS; 5 mM EDTA; 50 mM Na₂HPO₄; 250 µg/ml Herring sperm DNA; 50 µg/ml tRNA; or 0.25 M sodium phosphate buffer pH 7.2; 1 mM EDTA; 7% SDS

20

Hybridization temperature T = 65 to 70°C

25 Wash buffer: 0.2 × SSC; 0.1% SDS

Wash temperature T = 40 to 75°C.

- 30 Nucleic acid molecules which hybridize with the nucleic acid molecules according to the invention are capable, in principle, of encoding isoamylases from any wheat plant which expresses such proteins.

35 Nucleic acid molecules which hybridize with the molecules according to the invention can be isolated for example from genomic libraries or cDNA libraries of wheat or wheat plant tissue. Alternatively, they can be generated by recombinant methods or synthesized chemically.

Identification and isolation of such nucleic acid molecules can be effected using the molecules according to the invention or parts of these molecules or the reverse complements of these molecules, for example by means of

hybridization by standard methods (see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

- 5 Hybridization probes which can be used are, for example, nucleic acid molecules which have exactly or essentially the nucleotide sequence stated under Seq ID No. 1, 2 or 6 or parts of these sequences. The fragments used as hybridization probe may also be synthetic fragments which have been prepared with the aid of the customary synthetic techniques whose
10 sequence essentially agrees with that of a nucleic acid molecule according to the invention.

- The molecules hybridizing with the nucleic acid molecules according to the invention also encompass fragments, derivatives and allelic variants of the
15 above-described nucleic acid molecules which encode a wheat isoamylase according to the invention. Fragments are to be understood as meaning parts of the nucleic acid molecules of sufficient length so as to encode one of the proteins described. The term derivative means in this context that the sequences of these molecules differ from the sequences of the above-
20 described nucleic acid molecules at one or more positions and have a high degree of homology with these sequences. Homology means a sequence identity of at least 40%, in particular of at least 60%, preferably over 80%, especially preferably over 90%. The deviations relative to the above-described nucleic acid molecules may have been generated by deletion,
25 substitution, insertion or recombination.

- Homology furthermore means that functional and/or structural equivalence exists between the nucleic acid molecules in question or the proteins encoded by them. The nucleic acid molecules which are homologous to the
30 above-described molecules and constitute derivatives of these molecules are, as a rule, variations of these molecules which constitute modifications exerting the same biological function. They may be naturally occurring variations, for example, sequences from other organisms, or mutations which may have occurred naturally or been introduced by directed
35 mutagenesis. Furthermore, the variations may be synthetically generated sequences. The allelic variants may be both naturally occurring variants and synthetically generated variants or variants produced by recombinant DNA techniques.

The isoamylases encoded by the various variants of the nucleic acid molecules according to the invention share certain characteristics. These may include, for example, enzyme activity, molecular weight, immunological reactivity, conformation and the like, or else physical properties such as, for example, the migration behavior in gelelectrophoresis, the chromatographic behavior, sedimentation coefficients, solubility, spectroscopic characteristics, charge characteristics, stability; pH optimum, temperature optimum and the like.

10 The protein encoded by the nucleic acid molecules according to the invention is a wheat isoamylase. These proteins show certain homology ranges with isoamylases from other plant species which are already known.

The nucleic acid molecules according to the invention may be DNA molecules, in particular cDNA or genomic molecules. Furthermore, the nucleic acid molecules according to the invention may be RNA molecules which may result, for example, from the transcription of a nucleic acid molecule according to the invention. The nucleic acid molecules according to the invention may have been obtained, for example, from natural sources or they may have been generated by recombinant techniques or synthesized.

Subject matter of the invention are also oligonucleotides which hybridize specifically with a nucleic acid molecule according to the invention. Such oligonucleotides preferably have a length of at least 10, in particular of at least 15 and especially preferably of at least 50 nucleotides. The oligonucleotides according to the invention hybridize specifically with nucleic acid molecules according to the invention, i.e. not or only to a very low degree with nucleic acid sequences which encode other proteins, in particular other isoamylases. The oligonucleotides according to the invention can be used, for example, as primers for a PCR reaction or as hybridization probe for the isolation of the related genes. Equally, they may be constituents of antisense constructs or of DNA molecules encoding suitable ribozymes.

35 The invention furthermore relates to vectors, in particular plasmids, cosmids, phagemids, viruses, bacteriophages and other vectors conventionally used in genetic engineering comprising the above-described nucleic acid molecules according to the invention. Such vectors are

suitable for the transformation of pro- or eukaryotic cells, preferably plant cells.

5 The vectors especially preferably permit integration of the nucleic acid molecules according to the invention, if appropriate together with flanking regulatory regions, into the genome of the plant cell. Examples are binary vectors which can be employed in agrobacterial-mediated gene transfer. Preferably, integration of a nucleic acid molecule according to the invention in sense or antisense orientation ensures that a translatable or, if
10 appropriate, nontranslatable RNA is synthesized in the transformed pro- or eukaryotic cells.

The term "vector" generally denotes a suitable auxiliary known to the skilled worker which allows the directed transfer of a single- or double-stranded
15 nucleic acid molecule into a host cell, for example a DNA or RNA virus, a virus fragment, a plasmid construct which, in the absence or presence of regulatory elements, may be suitable for transferring nucleic acid into cells, or support materials such as glass fibers or else metal particles as can be employed, for example, in the particle gun method, but it may also
20 encompass a nucleic acid molecule which can be introduced directly into a cell by means of chemical or physical methods.

In a preferred embodiment, the nucleic acid molecules within the vectors are linked to regulatory elements which ensure transcription and synthesis
25 of a translatable RNA in pro- or eukaryotic cells or which – if desired – ensure synthesis of a nontranslatable RNA.

Expression of the nucleic acid molecules according to the invention in prokaryotic cells, for example, in *Escherichia coli*, is of importance for a
30 more detailed characterization of the enzymatic activities of the enzymes encoded by these molecules. In particular, it is possible to characterize the product synthesized by the enzymes in question in the absence of other enzymes involved in starch synthesis in the plant cell. This permits conclusions to be drawn regarding the function which the protein in
35 question exerts during starch synthesis in the plant cell.

In addition, various types of mutations can be introduced into the nucleic acid molecules according to the invention by means of customary techniques of molecular biology (see, for example, Sambrook et al., 1989,

Molecular Cloning, A Laboratory Manual, 2nd Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY), resulting in the synthesis of proteins whose biological properties may be altered. Possible here is, on the one hand, the generation of deletion mutants in which nucleic acid molecules are generated by successive deletions from the 5' or the 3' end of the coding DNA sequence which lead to the synthesis of correspondingly truncated proteins. Such deletions at the 5' end of the nucleotide sequence allow, for example, amino acid sequences to be identified which are responsible for translocation of the enzyme into the plastids (transit peptides). This allows the directed generation of enzymes which, owing to the removal of the sequences in question, are no longer localized in the plastids, but in the cytosol, or which, owing to the addition of other signal sequences, are localized in other compartments.

On the other hand, it is also possible to introduce point mutations at positions where an altered amino acid sequence affects, for example, enzyme activity or enzyme regulation. In this manner, it is possible to generate, for example, mutants which have an altered K_m value or which are no longer subject to the regulatory mechanisms via allosteric regulation or covalent modification which are normally present in the cell.

Furthermore, it is possible to generate mutants which have an altered substrate or product specificity of the protein according to the invention. Furthermore, it is possible to generate mutants which have an altered activity-temperature profile of the protein according to the invention.

To carry out the recombinant modification of prokaryotic cells, the nucleic acid molecules according to the invention or parts of these molecules can be introduced into plasmids which allow mutagenesis to take place or a sequence to be altered by recombining DNA sequences. Base exchanges can be carried out on natural or synthetic sequences added with the aid of standard methods (cf. Sambrook et al., 1989, Molecular Cloning: A laboratory manual, 2nd Ed., Cold Spring Harbor Laboratory Press, NY, USA). To link the DNA fragments to each other, adapters or linkers may be attached to the fragments. Furthermore, manipulations may be employed which provide suitable restriction cleavage sites or which eliminate superfluous DNA or restriction cleavage sites. Where insertions, deletions or substitutions are suitable, in vitro mutagenesis, primer repair, restriction or ligation may be employed. Analytical methods which are generally

employed are sequence analysis, restriction analysis or other methods of biochemistry and molecular biology.

5 In a further embodiment, the invention relates to host cells, in particular pro- or eukaryotic cells, which have been transformed with an above-described nucleic acid molecule according to the invention or a vector according to the invention, and to cells which are derived from cells transformed thus and comprise a nucleic acid molecule according to the invention or a vector. They are preferably pro- or eukaryotic cells, in
10 particular plant cells.

Subject matter of the invention are furthermore proteins with isoamylase activity which are encoded by the nucleic acid molecules according to the invention and which can be prepared by recombinant technology, and
15 processes for their preparation, where a host cell according to the invention is cultured under suitable conditions which are known to the skilled worker and which permit synthesis of the protein according to the invention and it is subsequently isolated from the host cells and/or the culture medium.

20 Providing the nucleic acid molecules according to the invention now makes it possible to intervene, with the aid of recombinant methods, in a directed fashion in the starch metabolism of plants and to alter it so that the resultant synthesis is of modified starch whose physicochemical properties, for example the amylose/amylopectin ratio, the degree of branching, the
25 average chain length, the phosphate content, the gelatinization behavior, the gel- or film-forming properties, the starch granule size and/or the starch granule shape is altered in comparison to known starch.

Thus, it is possible to express the nucleic acid molecules according to the
30 invention in plant cells in order to increase the activity of the isoamylase in question, or to introduce them into cells which do not naturally express this enzyme. Expressing the nucleic acid molecules according to the invention also makes it possible to lower the natural activity level of the isoamylase according to the invention in the plant cells. Furthermore, it is possible to
35 modify the nucleic acid molecules according to the invention by methods known to the skilled worker in order to obtain isoamylases according to the invention which are no longer subject to the cell's intrinsic regulatory mechanism or which have altered temperature-activity profiles or substrate or product specificities.

When expressing the nucleic acid molecules according to the invention in plants, it is possible, in principle, for the protein synthesized to be localized in any desired compartment of the plant cell. To achieve localization in a particular compartment, the sequence ensuring localization in plastids must be deleted and the remaining encoding region must, if necessary, be linked to DNA sequences which ensure localization in the compartment in question. Such sequences are known (see, for example, Braun et al., EMBO J. 11 (1992), 3219-3227; Wolter et al., Proc. Natl. Acad. Sci. USA 85 (1988), 846-850; Sonnewald et al., Plant J. 1 (1991), 95-106).

The present invention thus also relates to a method for generating transgenic plant cells which have been transformed with a nucleic acid molecule or a vector according to the invention, where a nucleic acid molecule according to the invention or a vector according to the invention is integrated into the genome of a plant cell, the transgenic plant cells which have been transformed by means of a vector or nucleic acid molecule according to the invention, and transgenic plant cells derived from cells transformed thus. The cells according to the invention comprise one or more nucleic acid molecules or vectors according to the invention, these preferably being linked to regulatory DNA elements which ensure transcription in plant cells, in particular to a suitable promoter. Such cells can be distinguished from naturally occurring plant cells inter alia by the fact that they comprise a nucleic acid molecule according to the invention which does not occur naturally in these cells, or by the fact that such a molecule exists integrated at a location in the cell's genome where it does not occur otherwise, i.e. in a different genomic environment. Furthermore, such transgenic plant cells according to the invention can be distinguished from naturally occurring plant cells by the fact that they comprise at least one copy of a nucleic acid molecule according to the invention stably integrated into the genome, if appropriate in addition to the copies of such a molecule which occur naturally in the cells. If the nucleic acid molecule(s) introduced into the cells is(are) additional copies to molecules which already occur naturally in the cells, then the plant cells according to the invention can be distinguished from naturally occurring plant cells in particular by the fact that this additional copy, or these additional copies, is, or are, localized at locations in the genome where it does not occur naturally, or they do not occur naturally. This can be checked, for example, with the aid of a Southern blot analysis.

If the nucleic acid molecule according to the invention which has been introduced into the plant genome is heterologous to the plant cell, the transgenic plant cells exhibit transcripts of the nucleic acid molecules according to the invention which can be detected in a simple manner by methods known to the skilled worker, for example by Northern blot analysis.

If the nucleic acid molecule according to the invention which has been introduced is homologous to the plant cell, the cells according to the invention can be distinguished from naturally occurring cells, for example, on the basis of the additional expression of nucleic acid molecules according to the invention. The transgenic plant cells preferably comprise more transcripts of the nucleic acid molecules according to the invention. This can be detected, for example, by Northern blot analysis. "More" in this context means preferably at least 10% more, preferably at least 20% more, especially preferably at least 50% more transcripts than corresponding untransformed cells. The cells furthermore preferably exhibit a corresponding increase or decrease in the activity of the protein according to the invention (at least 10%, 20% or 50%). The transgenic plant cells can be regenerated into intact plants by techniques known to the skilled worker.

Another subject matter of the present invention is a method for the generation of transgenic plants, where one or more nucleic acid molecules or vectors according to the invention are integrated into the genome of a plant cell and a complete plant is regenerated from said plant cell. Subject matter of the invention are furthermore plants which comprise the above-described transgenic plant cells. In principle, the transgenic plants can be plants of any species, i.e. not only monocotyledonous but also dicotyledonous plants. They are preferably useful plants, by preference starch-synthesizing or starch-storing plants, especially preferably rye, barley oats, wheat, sorghum and millet, sago, maize, rice, peas, marrowfat peas, cassava, potatoes, tomatoes, oilseed rape, soybeans, hemp, flax, sunflowers, cowpeas or arrowroot, in particular wheat, maize, rice and potatoes.

The invention also relates to propagation material of the plants according to the invention, for example fruits, seeds, tubers, rootstocks, seedlings, cuttings, calli, protoplasts, cell cultures and the like.

005077 27842960

The present invention furthermore relates to a process for the preparation of a modified starch comprising the step of extracting the starch from an above-described plant according to the invention and/or starch-storing parts of such a plant.

Processes for extracting the starch from plants or starch-storing parts of plants, in particular from wheat, are known to the skilled worker, cf., for example, Eckhoff et al. (Cereal Chem. 73 (1996) 54-57) "Starch: Chemistry and Technology (Eds.: Whistler, BeMiller and Paschall (1994), 2nd Edition, Academic Press Inc. London Ltd; ISBN 0-12-746270-8; see, for example, Chapter XII, pages 412-468: Corn and sorghum starches: production; by Watson; Chapter XIII, pages 469-479; Tapioca, arrowroot and sago starches: production; by Corbishley and Miller; Chapter XIV, pages 479-490: Potato starch: production and uses; by Mitch; Chapter XV, pages 491 to 506: Wheat starch: production, modification and uses; by Knight and Oson; and Chapter XVI, pages 507 to 528: Rice starch: production and uses; by Rohmer and Klem). Devices normally used in processes for extracting starch from plant material are separators, decanters, hydrocyclones, spray dryers and fluidized-bed dryers.

Owing to the expression of a nucleic acid molecule according to the invention, the transgenic plant cells and plants according to the invention synthesize a starch whose physicochemical properties, for example the amylose/amylopectin ratio, the degree of branching, the average chain length, the phosphate content, the gelatinization behavior, the starch granule size and/or starch granule shape is altered compared with starch synthesized in wild-type plants. In particular, such a starch may be altered with regard to viscosity and/or the film- or gel-forming properties of gels made from this starch in comparison with known starches.

Subject matter of the present invention is furthermore a starch which is obtainable from the plant cells and plants according to the invention and their propagation material and starch which is obtainable by the above-described process according to the invention.

It is furthermore possible to generate, with the aid of the nucleic acid molecules according to the invention, plant cells and plants in which the activity of a protein according to the invention is reduced. This also leads to

the synthesis of a starch with altered chemical and/or physical characteristics compared with starch from wild-type plant cells.

5 A further subject matter of the invention is thus also a transgenic plant cell comprising a nucleic acid molecule according to the invention in which the activity of an isoamylase is reduced in comparison with untransformed cells.

10 Plant cells with a reduced activity of an isoamylase can be obtained, for example, by expressing a suitable antisense RNA, a sense RNA for achieving a cosuppression effect or by expressing a suitably constructed ribozyme which specifically cleaves transcripts which encode an isoamylase, making use of the nucleic acid molecules according to the invention by methods known to the skilled worker, cf. Jorgensen (Trends
15 Biotechnol. 8 (1990), 340-344), Niebel et al., (Curr. Top. Microbiol. Immunol. 197 (1995), 91-103), Flavell et al. (Curr. Top. Microbiol. Immunol. 197 (1995), 43-46), Palaqui and Vaucheret (Plant. Mol. Biol. 29 (1995), 149-159), Vaucheret et al., (Mol. Gen. Genet. 248 (1995), 311-317), de Borne et al. (Mol. Gen. Genet. 243 (1994), 613-621).

20 To reduce the activity of an isoamylase according to the invention, it is preferred to reduce, in the plant cells, the number of transcripts encoding it, for example by expressing an antisense RNA.

25 Here, it is possible to make use, on the one hand, of a DNA molecule which encompasses all of the sequence encoding a protein according to the invention, inclusive of any flanking sequences which may be present, or else of DNA molecules which only encompass parts of the coding sequence, it being necessary for these parts to be sufficiently long so as to
30 cause an antisense effect in the cells. In general, sequences up to a minimum length of 15 bp, preferably with a length of 100-500 bp, may be used for efficient antisense inhibition in particular sequences with a length of over 500 bp. As a rule, DNA molecules are used which are shorter than 5000 bp, preferably sequences which are shorter than 2500 bp.

35 Also possible is the use of DNA sequences which show a high degree of homology with the sequences of the DNA molecules according to the invention, but are not completely identical. The minimum homology should

exceed approx. 65%. The use of sequences with homologies between 95 and 100% is to be preferred.

5 Subject matter of the invention is also a process for producing a modified starch encompassing the step of extracting the starch from a cell or plant according to the invention and/or from starch-storing parts of such a plant.

10 Subject matter of the invention is furthermore starch which can be obtained from the cells or plants according to the invention and their propagation material or parts, and also starch which can be obtained by a process according to the invention.

15 The starches according to the invention can be modified by methods known to the skilled worker and are suitable, in their unmodified or modified form, for a variety of applications in the food or non-food sectors.

20 In principle, the possible uses of the starches according to the invention can be divided into two important sectors. One sector encompasses the hydrolyzates of the starch, mainly glucose and glucan units, which are obtained by enzymatic or chemical methods. They are used as starting material for further chemical modifications and processes such as fermentation. What would be feasible for reducing the costs is the simplicity and economic design of a hydrolytic method. It currently proceeds essentially enzymatically using amyloglucosidase. What would be feasible
25 is a financial saving by using less enzyme. This could be caused by altering the structure of the starch, for example by increasing the surface area of the granule, better digestibility, for example owing to a lower degree of branching or a sterical structure which limits the accessibility for the enzymes employed.

30

The other sector in which the starch according to the invention can be used as so-called native starch, owing to its polymeric structure, can be divided into two further fields of application:

35 1. The food industry

Starch is a traditional additive to a large number of foodstuffs in which its function is essentially to bind aqueous additives or to cause increased viscosity or else increased gelling. Important characteristics are the rheology, the sorptive characteristics, the

swelling temperature, the gelatinization temperature, the viscosity, the thickening power, the starch solubility, the transparency and gel structure, the thermal stability, the shear stability, the stability to acids, the tendency to undergo retrogradation, the film-forming capacity, the freeze-thaw stability, the viscosity stability in salt solutions, the digestibility and the ability to form complexes with, for example, inorganic or organic ions.

2. The non-food industry

In this large sector, starch can be employed as auxiliary for various preparation processes or as an additive in industrial products. When using starch as an auxiliary, mention must be made, in particular, of the paper and board industry. Starch acts mainly for retardation purposes (retaining solids), for binding filler particles and fines, as stiffener and for dehydration. Moreover, the advantageous properties of starch regarding stiffness, strength, sound, touch, luster, smoothness, bonding strength and the surfaces is utilized.

2.1 Paper and board industry

Within the papermaking process, four fields of application must be distinguished, i.e. surface, coating, stock and spraying. The demands on starch with regard to surface treatment are essentially high whiteness, an adapted viscosity, high viscosity stability, good film formation and low dust formation. When used for coating, the solids content, a suitable viscosity, a high binding capacity and a high pigment affinity play an important role. Of importance when used as additive to the stock is rapid, uniform, loss-free distribution, high mechanical strength and complete retention in the paper web. If the starch is used in the spraying sector, again, an adapted solids content, high viscosity and high binding capacity are of importance.

2.2 The adhesives industry

An important field of application for starches is the adhesives industry, where the potential uses can be divided into four subsections: the use as a pure starch paste, the use in starch pastes which have been treated with specialty chemicals, the use of starch as additive to synthetic resins and polymer dispersions, and the use of starches as extenders for synthetic adhesives. 90% of the starch-based adhesives are employed in the sectors production of

corrugated board, production of paper sacks and bags, production of composite materials for paper and aluminum, production of boxes and gumming adhesives for envelopes, stamps and the like.

2.3 Textile industry and textile care products industry

5 An important field of application for starches as auxiliaries and additives is the sector production of textiles and textile care products. The following four fields of application must be distinguished within the textile industry: the use of starch as sizing agent, i.e. as auxiliary for smoothing and strengthening the
10 smoothing behavior as protection from the tensile forces applied during weaving, and for increasing abrasion resistance during weaving, starch as a textile finishing agent, in particular after quality-reducing pretreatments such as bleaching, dyeing and the like, starch as thickener in the preparation of dye pastes for preventing
15 bleeding, and starch as additive to glazing agents for sewing threads.

2.4 Construction materials industry

20 The fourth field of application is the use of starches as additives in construction materials. An example is the production of gypsum plasterboards, where the starch which is admixed to the gypsum slurry gelatinizes with the water, diffuses to the surface of the plaster core and there binds the board to the core. Other fields of application are the admixture to rendering and mineral fibers. In the
25 case of ready-mixed concrete, starch products are employed for delaying binding.

2.5 Soil stabilization

30 Another market for starch products is the production of soil stabilizers, which are employed for the temporary protection of the soil particles from water when the soil is disturbed artificially. According to present knowledge, product combinations of starch and polymer emulsions equal the previously employed products with regard to their erosion- and crust-reducing effect, but are markedly
35 less expensive.

2.6 Use in crop protection products and fertilizers

One field of application for using starch is in crop protection products for altering the specific properties of the products. Thus, starch can

be employed for improving the wettability of crop protection products and fertilizers, for the controlled release of the active ingredients, for converting liquid, volatile and/or malodorous active ingredients into microcrystalline, stable, shapeable substances, for mixing incompatible compounds and for extending the duration of action by reducing decomposition.

2.7 Pharmaceuticals, medicine and cosmetics industry

Another field of application is the sector of the pharmaceuticals, medicine and cosmetics industry. In the pharmaceuticals industry, starch can be employed as binder for tablets or for diluting the binder in capsules. Moreover, starch can be employed as tablet disintegrant since it absorbs fluid after swallowing and swells within a short time to such an extent that the active ingredient is liberated. Medicinal lubricating powders and wound powders are starch-based for reasons of quality. In the cosmetics sector, starches are employed, for example, as carriers of powder additives such as fragrances and salicylic acid. A relatively large field of application for starch is toothpaste.

2.8 Addition of starch to coal and briquettes

A field of application for starch is as additive to coal and briquettes. With an addition of starch, coal can be agglomerated, or briquetted, in terms of high quantity, thus preventing premature decomposition of the briquettes. In case of barbecue coal, the starch addition amounts to between 4 and 6%, in the case of calorized coal to between 0.1 and 0.5%. Moreover, starches are gaining importance as binders since the emission of noxious substances can be markedly reduced when starches are added to coal and briquettes.

2.9 Ore slick and coal silt processing

Furthermore, starch can be employed as flocculant in the ore slick and coal silt processing sector.

2.10 Foundry auxiliary

A further field of application is as additive to foundry auxiliaries. Various casting processes require cores made with sands treated with binders. The binder which is predominantly employed

water vapor, an improved antistatic behavior, an improved antiblock behavior and an improved printability with aqueous inks can be achieved.

Another possibility is the use of starch in polyurethane foams. By adapting the starch derivatives and by process-engineering optimization, it is possible to control the reaction between synthetic polymers and the starches' hydroxyl groups in a directed manner. This results in polyurethane films which have the following spectrum of properties, owing to the use of starch: a reduced heat expansion coefficient, a reduced shrinking behavior, an improved pressure-tension behavior, an increase in permeability for water vapor without altering the uptake of water, a reduced flammability and a reduced ultimate tensile strength, no drop formation of combustible parts, freedom from halogens, or else reduced aging. Disadvantages which still exist are reduced printability and reduced impact strength.

Product development is currently no longer restricted to films. Solid polymer products such as pots, slabs and dishes which have a starch content of over 50% may also be produced. Moreover, starch/polymer mixtures are considered advantageous since their biodegradability is much higher.

Starch graft polymers have become exceedingly important owing to their extremely high water binding capacity. They are products with a starch backbone and a side lattice of a synthetic monomer, grafted on following the principle of the free-radical chain mechanism. The starch graft polymers which are currently available are distinguished by a better binding and retention capacity of up to 1000 g of water per g of starch combined with high viscosity. The fields of application of these superabsorbers have extended greatly in recent years and are, in the hygiene sector, products such as diapers and pads and, in the agricultural sector, for example seed coatings.

What is decisive for the application of novel, genetically modified starches are, on the one hand, structure, water content, protein content, lipid content, fiber content, ash/phosphate content, amylose/amylopectin ratio, molecular mass distribution, degree of branching, granule size and granule shape and crystallinity, and, on the other hand, also the characteristics which effect the following features: flow and sorption behavior,

gelatinization temperature, viscosity, viscosity stability in salt solutions, thickening power, solubility power, gel structure and gel transparency, thermal stability, shear stability, stability to acids, tendency to undergo retrogradation, gel formation, freeze-thaw stability, complex formation, iodine binding, film formation, adhesive power, enzyme stability, digestibility and reactivity.

The production of modified starches by recombinant methods can, on the one hand, alter the properties of the starch derived from the plant in such a way that other modifications by means of chemical or physical processes are no longer required. On the other hand, starches which have been altered by recombinant methods may be subjected to further chemical modification, which leads to further improvements in quality for some of the above-described fields of application. These chemical modifications are known in principle. They are, in particular, modifications by thermal treatment, treatment with organic or inorganic acids, oxidation and esterifications, which lead, for example, to the formation of phosphate starches, nitrate starches, sulfate starches, xanthate starches, acetate starches and citrate starches. Moreover, mono- or polyhydric alcohols in the presence of strong acids may be employed for producing starch ethers, resulting in starch alkyl ethers, O-allyl ethers, hydroxyalkyl ethers, O-carboxy methyl ethers, N-containing starch ethers, P-containing starch ethers), S-containing starch ethers, crosslinked starches or starch graft polymers.

A preferred use of the starches according to the invention is the production of packaging materials and disposable articles, on the one hand, and as foodstuff or foodstuff precursor on the other hand.

To express the nucleic acid molecules according to the invention in sense or antisense orientation in plant cells, they are linked to regulatory DNA elements which ensure transcription in plant cells. These include, in particular, promoters, enhancers and terminators. In general, any promoter which is active in plant cells is suitable for expression.

The promoter may be chosen in such a way that expression is constitutive or takes place only in a particular tissue, at a particular point in time of plant development or at a point in time determined by external factors. Relative to the plant, the promoter can be homologous or heterologous. Examples

of suitable promoters are the cauliflower mosaic virus 35S RNA promoter and the maize ubiquitin promoter for constitutive expression, the patatin promoter B33 (Rocha-Sosa et al., EMBO J. 8 (1989), 23-29) for tuber-specific expression, or a promoter which ensures expression only in photosynthetically active tissue, for example the ST-LS1 promoter (Stockhaus et al., Proc. Natl. Acad. Sci. USA 84 (1987), 7943-7947; Stockhaus et al., EMBO J. 8 (1989), 2445-2451) or, for endosperm-specific expression, the wheat AMG promoter, the USP promoter, the phaseolin promoter or promoters from maize zein genes.

10

A termination sequence which serves to correctly terminate transcription and to add a poly-A tail to the transcript, which is considered to have a function in stabilizing the transcripts, may also be present. Such elements have been described in the literature (cf. Gielen et al., EMBO J. 8 (1989), 23-29) and are exchangeable as desired.

15

The present invention provides nucleic acid molecules which encode a protein with a wheat isoamylase function. The nucleic acid molecules according to the invention permit the production of this enzyme whose functional identification in starch biosynthesis, the generation of plants which have been altered by recombinant technology in which the activity of this enzyme is altered and thus allows a starch to be synthesized in plants modified thus whose structure is altered and whose physicochemical properties are altered.

20

In principle, the nucleic acid molecules according to the invention may also be used for generating plants in which the activity of the isoamylase according to the invention is increased or reduced while simultaneously the activities of other enzymes which participate in starch synthesis are altered.

25

Altering the activities of an isoamylase in plants results in the synthesis of a starch with altered structure. Furthermore, nucleic acid molecules which encode an isoamylase or suitable antisense constructs can be introduced into plant cells in which the synthesis of endogenous starch synthases or branching enzymes is already inhibited (as, for example, in WO 92/14827 or Shannon and Garwood, 1984, in Whistler, BeMiller and Paschall, Starch: Chemistry and Technology, Academic Press, London, 2nd Edition: 25-86).

30

If it is intended to achieve the inhibition of the synthesis of several enzymes involved in starch biosynthesis in transformed plants, the transformation

may involve DNA molecules which simultaneously comprise several regions encoding the enzymes in question in antisense orientation under the control of a suitable promoter. Here, it is possible for each sequence to be under the control of its own promoter, or for the sequences to be transcribed by a joint promoter as a fusion or to be under the control of a joint promoter. The last-mentioned alternative will generally be preferred, since in this case the synthesis of the proteins in question should be inhibited roughly to the same extent. As regards the length of the individual coding regions used in such a construct, what has been mentioned above for the generation of antisense constructs also applies here. In principle, there is no upper limit for the number of antisense fragments to transcribed in such a DNA molecule starting from one promoter. However, the transcript formed should preferably not exceed a length of 10 kb, in particular a length of 5 kb.

Coding regions localized in such DNA molecules in combination with other coding regions in antisense orientation behind a suitable promoter may be derived from DNA sequences which encode the following proteins: starch-granule-bound starch synthases (GBSS I and II) and soluble starch synthases (SSS I and II), branching enzymes (isoamylases, pullulanases, R enzymes, branching enzymes, debranching enzymes), starch phosphorylases and disproportioning enzymes. This enumeration is only by way of example. The use of other DNA sequences for the purposes of such a combination is also feasible.

Such constructs allow the synthesis of a plurality of enzymes to be inhibited simultaneously in plant cells transformed with them.

Furthermore, the constructs can be introduced into plant mutants which are deficient for one or more starch biosynthesis genes (Shannon and Garwood, 1984, in Whistler, BeMiller and Paschall, Starch: Chemistry and Technology, Academic Press, London, 2nd Edition: 25-86). These defects may relate to the following proteins: starch-granule-bound starch synthases (GBSS I and II) and soluble starch synthases (SSS I and II), branching enzymes (BE I and II), debranching enzymes (R enzymes), disproportioning enzymes and starch phosphorylases. This enumeration is only by way of example.

Such a procedure furthermore allows the synthesis of a plurality of enzymes to be inhibited simultaneously in plant cells transformed with them.

- 5 To prepare the introduction of foreign genes into higher plants, a large number of cloning vectors containing a replication signal for E.coli and a marker gene for selecting transformed bacterial cells is available. Examples of such vectors are pBR322, pUC series, M13mp series, pACYC184 and the like. The desired sequence may be introduced into the vector at a
- 10 suitable restriction cleavage site. The plasmid obtained is used to transform E.coli cells. Transformed E.coli cells are grown in a suitable medium and subsequently harvested and lyzed. The plasmid is recovered. Analytical methods for characterizing the plasmid DNA obtained which are generally used are restriction analyses, gel electrophoresis and further methods of
- 15 biochemistry and molecular biology. After each manipulation, the plasmid DNA can be cleaved and resulting DNA fragments linked to other DNA sequences. Each plasmid DNA sequence can be cloned in identical or different plasmids.
- 20 A large number of techniques is available for introducing DNA into a plant host cell. These techniques encompass transformation of plant cells with t-DNA using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as transformation agents, protoplast fusion, injection, the electroporation of DNA, the introduction of DNA by means of the biolistic method, and other
- 25 possibilities.

- The injection and electroporation of DNA into plant cells per se require no particular aspect of the plasmids used. Simple plasmids such as, for example, pUC derivatives may be used. However, if intact plants are to be
- 30 regenerated from such transformed cells, the presence of a selectable marker gene is generally required.

- Depending on the method of introducing desired genes into the plant cell, further DNA sequences may be required. If, for example, the Ti or Ri
- 35 plasmid is used for transforming the plant cell, at least the right border, but frequently the right and left borders, of the Ti and Ri plasmid T-DNA must be linked to the genes to be introduced as flanking region.

If agrobacteria are used for the transformation, the DNA to be introduced must be cloned into specific plasmids, either into an intermediate vector or into a binary vector. The intermediate vectors can be integrated into the agrobacterial Ti or Ri plasmid by homologous recombination owing to sequences which are homologous to sequences in the T-DNA. The former also contains the vir region, which is required for the T-DNA transfer. Intermediate vectors cannot replicate in agrobacteria. The intermediate vector can be transferred to *Agrobacterium tumefaciens* (conjugation) by means of a helper plasmid. Binary vectors are capable of replication in *E. coli* and in agrobacteria. They contain a selection marker gene and a linker or polylinker, which are framed by the left and right T-DNA border regions. They can be transformed directly into the agrobacteria (Holsters et al. *Mol. Gen. Genet.* 163 (1978), 181-187). The agrobacterium which acts as the host cell should contain a plasmid carrying a vir region. The vir region is required for transferring the T-DNA into the plant cell. Additional T-DNA may be present. The agrobacterium thus transformed can be used for transforming plant cells.

The use of T-DNA for transforming plant cells has been researched intensively and been described in EP 120 516; Hoekema, In: *The Binary Plant Vector System* Offsetdrukkerij Kanters B.V., Alblasterdam (1985), Chapter V; Fraley et al., *Crit. Rev. Plant. Sci.*, 4, 1-46 and An et al. *EMBO J.* 4 (1985), 277-287.

To transfer the DNA into the plant cell, plant explants can expediently be cocultured with *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. Intact plants can then be regenerated again from the infected plant material (for example leaf sections, stalk sections, roots, but also protoplasts, or plant cells grown in suspension culture) in a suitable medium which can contain, inter alia, certain sugars, amino acids, antibiotics or biocides for selecting transformed cells. The resulting plants can then be examined for the presence of the DNA which has been introduced. Other possibilities of introducing foreign DNA using the biolistic method or by protoplast transformation are known (cf., for example, Willmitzer, L., 1993 *Transgenic plants*. In: *Biotechnology, A Multi-Volume Comprehensive Treatise* (H.J. Rehm, G. Reed, A. Pühler, P. Stadler, eds.), Vol. 2, 627-659, VCH Weinheim-New York-Basel-Cambridge).

While the transformation of the dicotyledonous plants via Ti-plasmid vector systems with the aid of *Agrobacterium tumefaciens* is well established, more recent work suggests that even monocotyledonous plants are indeed accessible to transformation by means of agrobacterium-based vectors
 5 (Chan et al., *Plant Mol. Biol.* 22 (1993), 491-506; Hiei et al., *Plant J.* 6 (1994), 271-282).

Alternative methods for the transformation of monocotyledonous plants are the transformation by means of the biolistic approach, protoplast
 10 transformation, or the physically- or chemically-induced DNA uptake into protoplasts, for example by electroporation of partially permeabilized cells, transfer of DNA by means of glass fibers, macroinjection of DNA into inflorescences, the microinjection of DNA into microspores or proembryos, DNA uptake by germinating pollen and DNA uptake in embryos by swelling
 15 (review: Potrykus, *Physiol. Plant* (1990), 269-273).

Three of the abovementioned transformation systems have been established in the past for various cereals: the electroporation of tissue, the transformation of protoplasts and the DNA transfer by particle
 20 bombardment into regenerable tissue and cells (review: Jähne et al., *Euphytica* 85 (1995), 35-44).

Different methods of transforming wheat have been described in the literature (review: Maheshwari et al., *Critical Reviews in Plant Science* 14
 25 (2) (1995), 149 to 178); Hess et al. (*Plant Sci.* 72 (1990), 233) employ the macroinjection method to bring pollen and agrobacteria into the immediate vicinity. The mobilization of the plasmid which contained the *nptII* gene as selectable marker was detected by Southern blot analysis and NPTII test. The transformants showed a normal phenotype and were fertile.
 30 Kanamycin resistance was detected in two consecutive generations.

The first transgenic fertile wheat plant which was regenerated after bombardment with DNA bound to microprojectiles was described by Vasil et al. (*Bio/Technology* 10 (1992), 667-674). The target tissue for the
 35 bombardment was an embryogenic callus culture (type C callus). The selection marker employed was the *bar* gene which encodes a phosphinothricin acetyl transferase and thus mediates resistance to the herbicide phosphinothricin. A further system was described by Weeks et al. (*Plant Physiol.* 102 (1993), 1077-1084), and Becker et al. (*Plant J.* 5(2)

(1994), 299-307). Here, the target tissue for the DNA transformation is the scutellum of immature embryos which was stimulated in a preliminary in-vitro phase to induce somatic embryos. The transformation efficacy in the system developed by Becker et al. (loc cit.) is 1 transgenic plant per 83 embryos of the variety "Florida" and thus markedly higher than the system established by Weeks et al., which yields 1 to 2 transgenic plants per 1000 embryos of the variety "Bohwhite".

The system developed by Becker et al. (loc Cit) forms the basis for the transformation experiments described in the examples.

10

Once the DNA introduced is integrated into the genome of the plant cell, it is, as a rule, stable and is also retained in the progeny of the originally transformed cell. It normally contains one of the above-mentioned selection markers which mediates, for example, resistance to a biocide such as phosphinothricin or an antibiotic such as kanamycin, G 418, bleomycin or hygromycin, to the transformed plant cells or which permits selection via the presence or absence of certain sugars or amino acids. The marker chosen individually should therefore allow the selection of transformed cells over cells which lack the DNA introduced.

20

Within the plant, the transformed cells grow in the customary manner (see also McCormick et al., Plant Cell Reports 5 (1986), 81-84). The resulting plants can be grown normally and hybridized with plants which have the same transformed germ plasm or other germ plasm. The resulting hybrid individuals have the corresponding phenotype properties. Seeds may be obtained from the plant cells. Two or more generations should be grown in order to ensure that the phenotype characteristic is stably retained and inherited. Also, seeds should be harvested in order to ensure that the phenotype in question or other characteristics have been retained.

30

The examples which follow are intended to illustrate the invention and do not constitute any restriction whatsoever.

1. Cloning methods

35

The vector pBluescript II SK (Stratagene) was used for cloning in E.coli.

2. Bacterial strains

The *E. coli* strain DH5 α (Bethesda Research Laboratories, Gaithersburg, USA) was used for the Bluescript vector and for the antisense constructs. The *E. coli* strain XL1-Blue was used for the in vivo excision.

3. Transformation of immature wheat embryos

Media

MS:	100 ml/l macrosalt 1 ml/l microsalt 2 ml/l Fe/NaEDTA 30 g/l sucrose	(D. Becker and H. Lörz, Plant Tissue Culture Manual (1996), B 12:1-20)
#30:	MS + 2,4-D (2 mg/l)	
#31:	MS + 2,4-D (2 mg/l) + phosphinothricin (PPT, active component of herbicide BASTA (2 mg/l))	
#32:	MS + 2,4-D (0.1 mg/l) + PPT (2 mg/l)	
#39:	MS + 2,4-D (2 mg/ml) + of each 0.5 N mannitol/sorbitol	

The media stated were brought to pH 5.6 using KOH and solidified using 0.3% Gelrite.

The method for transforming immature wheat embryos was developed and optimized by Becker and Lörz (D. Becker and H. Lörz, Plant Tissue Culture Manual (1996), B12: 1 to 20).

In the experiments described hereinbelow, the procedure developed by Becker and Lörz (loc. Cit) was adhered to.

For the transformation, ears with caryopses of developmental stage 12 to 14 days after anthesis were harvested and surface-sterilized. The isolated scutella were plated onto induction medium #30 with the embryo axis orientated toward the medium.

After preculture for 2 to 4 days (26°C, in the dark), the explants are transferred to medium #39 for the osmotic preculture (2 to 4 h, 26°C, in the dark).

- 5 For the biolistic transformation, approx. 29 µg of gold particles on which a few µg of the target DNA had previously been precipitated were employed per shot. Since the experiments carried out are cotransformants, the target DNA composed of the target gene and a resistance marker gene (bar gene) in the ratio 1:1 is added to the precipitation batch.

10

4. DIG labeling of DNA fragments

- DNA fragments employed as screening probes were labeled via a specific PCR with the incorporation of DIG-labeled dUTP (Boehringer Mannheim, Germany).

15

Media and solutions used in the examples:

- | | |
|----------|---|
| 20 × SSC | 175.3 g NaCl |
| 20 | 88.2 g sodium citrate |
| | twice-distilled H ₂ O to 1000 ml |
| | 10 N NaOH to pH 7.0 |

- Plasmid pTaSU 8A was deposited at the DSMZ in Braunschweig, Federal Republic of Germany, as specified in the Budapest Treaty under the No. DSM 12795, and plasmid pTaSU 19 under the No. DSM 12796.

25

- Example 1: Identification, isolation and characterization of a cDNA encoding an isoamylase ("sugary" homolog) from wheat (*Triticum aestivum* L., cv Florida)

30

To identify a cDNA which encodes a wheat isoamylase isoform (sugary), a heterologous screening strategy was followed. To this end, a wheat cDNA library was screened with a maize sugary probe.

35

The probe (sugary probe) was isolated from a maize cDNA library by means of specific primers using PCR amplification. The maize cDNA library was cloned from poly(A) + RNA from a mixture of equal amounts of 13-, 17-, 19-, 20-, 22-, 25- and 28- day (DAP) old caryopses in a Lambda Zap II

vector following the manufacturer's instructions (Lambda ZAP II-cDNA Synthesis Kit Stratagene GmbH, Heidelberg, Germany). In all the caryopses used, with the exception of the 13-day-old kernels, the embryo had been removed prior to isolating the RNA.

5

The DNA fragment employed as probe for screening the wheat cDNA library was amplified with the following primers:

su1p-1: 5'AAAGGCCCAATATTATCCTTTAGG 3' (Seq ID No. 4)

su1p-2: 5' GCCATTTCAACCGTTCTGAAGTCGGAAGTC 3' (Seq. ID No. 5)

10

The template employed for the PCR reaction was 2 µl of the amplified maize cDNA library. Furthermore, the PCR reaction contained 1.5-3 mM MgCl₂, 20 mM Tris-HCl (pH 8.4), 50 mM KCl, 0.8 mM dNTP mix, 1 µM primer su1p-1a, 1 µM primer su1p-2 and 2.5 units Taq polymerase (recombinant, Life Technologies).

15

The amplification was carried out using a Trioblock from Biometra following the scheme: 4 min/95°C; 1 min/95°C, 45 sec/58°C; 1 min 15 sec/72°C; 30 cycles 5 min/72°C. The amplified DNA band of approx. 990 bp was separated in an agarose gel and excised. A second amplification was [lacuna] from this fragment following the scheme as described above. The 990 bp fragment obtained from this second amplification was cleaved with the restriction enzyme *BAM* HI into a 220 bp and a 770 bp fragment. After the sugary fragment had again been separated in an agarose gel, the band excised and the fragment isolated, the probe was DIG-labeled. 500 ng of sugary fragment were employed for the random-prime labeling with digoxigenin. 10 µl of random primer were added to the fragment to be labeled and the reaction was heated for 5 min at 95-100°C. After heating, 0.1 mM dATP, 0.1 mM dGTP, 0.1 mM dCTP and 0.065 mM dTTP and 0.035 mM digoxigenin-11-dUTP (Boehringer Mannheim) and Klenow buffer (standard) and 1 unit of Klenow polymerase were added. The reaction was allowed to proceed at RT (room temperature) overnight. To check the labeling, a dot test was carried out following the manufacturer's instructions ("The DIG System User's Guide for Filter Hybridization" by

25

30

35

The wheat cDNA library was synthesized from poly(A) + RNA of approx. 21-day ("starchy" endosperm) old caryopses in a Lambda Zap II vector following the manufacturer's instructions (Lambda ZAP II-cDNA Synthesis

Kit, Stratagene GmbH, Heidelberg). After determination of the titer of the cDNA library, a primary titer of 1.26×10^6 pfu/ml was determined.

To screen the wheat cDNA library, approx. 350,000 phages were plated out. The phages were plated out and the plates blotted by standard protocols. The filters were prehybridized and hybridized in 5X SSC, 3% blocking (Boehringer, Mannheim), 0.2% SDS, 0.1% sodium laurylsarcosine and 50 µg/ml herring sperm DNA at 55°C. 1 ng/ml of the labeled sugary probe was added to the hybridization solution and the hybridization was incubated overnight. The filters were washed 2 x 5 mins in 2 X SSC, 1% SDS at RT; 2 x 10 min in 1 X SSC, 0.5% SDS at 55°C; 2 x 10 min in 0.5 X SSC, 0.2% SDS at 55°C. Positive clones were singled out by further screening cycles. Single clones were obtained via in vivo excision as pBluescript SK phagemids (procedure analogous to the manufacturer's instructions; Stratagene, Heidelberg, Germany).

After the clones had been analyzed via minipreps and restriction of the plasmid DNA, clone pTaSU-19 was deposited at the DSMZ Deutsche Sammlung für Mikroorganismen und Zellkulturen GmbH under the number DSM 12796 and analyzed in greater detail.

Example 2: Sequence analysis of cDNA insertions of plasmids pTaSU19

The plasmid DNA was isolated from clone pTaSU19 and the sequence of the cDNA insertions determined by means of the dideoxynucleotide method (Sanger et al., Proc. Natl. Acad. Sci. USA 74 (1977), 5463-5467).

The insertion of clone TaSU-19 is 2997 bp in length and constitutes a partial cDNA. The nucleotide sequence is shown under Seq ID No. 1. A comparison with already published sequences revealed that the sequence shown under Seq ID No. 1 encompasses a coding region which has homologies to isoamylases from other organisms.

Sequence analysis also reveals that two introns are located in the cDNA sequence in positions 297-396 (intron 1) and 1618-2144 (intron 2). If these introns are removed, a protein sequence may be derived which exhibits homologies to the protein sequences of isoamylases of other organisms. The amino acid sequence which corresponds to the coding regions of Seq ID No. 1 is shown under Seq ID No. 3.

Example 3 Generation of the plant transformation vector pTa-alpha-SU19

To express an antisense RNA corresponding to the TaSU19-cDNA, the
 5 plant transformation vectors pTa-alpha-SU19 were constructed on the
 basis of the basic plasmid pUC19 by linking the cDNA insertion of plasmid
 pTa-alpha-SU19 in antisense orientation to the 3' end of the ubiquitin
 promoter. This promoter is composed of the first untranslated exon and the
 first intron of the maize *ubiquitin 1* gene (Christensen A.H. et al., Plant
 10 Molecular Biology 19 (1992), 675-689). Parts of the polylinker and the NOS
 terminator are derived from plasmid pAct1.cas (CAMBIA, TG 0063;
 Cambia, GPO Box 3200, Canberra ACT 2601, Australia). Vector constructs
 with this terminator and constructs based on pAct1.cas are described by
 MCElroy et al. (Molecular Breeding 1 (1995), 27-37). The vector thus
 15 formed was termed pUbi.cas.

The vector was cloned by restricting a 2kb fragment from clone Ta-SU19
 with the restriction enzyme *Xba I*. The fragment was filled up at the ends by
 means of Klenow reaction and subsequently ligated into the *Sma I* cloning
 20 site of the expression vector pUbi.cas.

The resulting expression vector is termed Ta-alpha-SU 19 and is used as
 described above for transforming wheat.

25 Example 4: Isolation and characterization of a further cDNA encoding an
 isoamylase (sugary 1 homolog) from wheat (*Triticum aestivum* L., cv
 Florida)

A wheat cDNA library was screened with a sugary probe which represents
 30 a part of clone pTaSU19, viz. positions 489-1041 of Seq ID No. 1.

The wheat-specific digoxigenin-labeled sugary probe employed for
 screening the cDNA library was prepared by means of PCR amplification.
 The primers employed in this reaction were:

35 SUSO1: 5'-GCT TTA CGG GTA CAG GTT CG-3' (Seq ID No. 8), and
 SUSO2: 5'-AAT TCC CCG TTT GTG AGC-3' (Seq ID No. 9)

1 ng of plasmid pTaSU19 was employed in the reaction as template. In
 addition, the PCR reaction contained in each case 300 nM of the primers

SUSO1 and SUSO2, in each case 100 μ M of the nucleotides dATP, dGTP, dCTP, 65 μ M dTTP, 35 μ M digoxigenin-11-dUTP (Boehringer Mannheim), 1.5 mM $MgCl_2$, and 2.5 U (units) Taq polymerase and 10 μ l of 10-fold concentrated Taq polymerase reaction buffer (both Life Technologies). The final volume of the reaction was 100 μ l. The amplification was performed in a PCR apparatus (TRIO[®] Thermoblock, Biometra) with the following temperature regime: 3 min at 95°C (once); 45 sec at 95°C – 45 sec at 55°C – 2 min at 72°C (30 cycles); 5 min at 72°C (once). A 553 bp DNA fragment was revealed owing to the reduced mobility in the agarose gel in comparison with the product of a controlled reaction without digoxigenin-11-dutp.

The caryopses-specific wheat cDNA library of Example 1 was screened with the resulting digoxigenin-labeled probe.

The hybridization step was performed overnight in 5x SSC, 0.2% SDS, 0.1% sodium laurylsarcosin and 50 μ g/ml herring sperm DNA at 68°C in the presence of 1 ng/ml of the digoxigenin-labeled probe. After the hybridization, the filters were washed as follows: 2 x 5 min in 2x SSC, 1% SDS at RT; 2 x 10 min in 1x SSC, 0.5% SDS at 68°C; 2 x 10 min in 0.5x SSC, 0.2% SDS at 68°C. Positive clones were singled out by at least two further screening cycles. Plasmids were obtained from the phage clones pBluescript SK via *in vivo* excision (protocols in accordance with the manufacturer's instructions; Stratagene, Heidelberg, Germany). After restriction analysis it clones obtained, clone pTaSU8A was deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen under the number DSM 12795 and studied in greater detail.

Example 5: Sequence analysis of the cDNA insert in plasmid pTaSU8A

The nucleotide sequence of the cDNA insert in plasmid pTaSU8A was determined by means of the dideoxynucleotide method (Seq ID No. 6).

The insertion of clone pTaSU8A is 2437 bp in length and constitutes a partial cDNA. A comparison with already published sequences reveals that the sequence shown under Seq ID No. 6 comprises a coding region which has homologies to isoamylases from other organisms. Equally, the protein sequence derived from the coding region of clone pTaSU8A and shown in

Seq ID No. 7 exhibits homologies to the protein sequences of isoamylases of other organisms. Upon comparison of the sequences of clones pTaSU19 (Seq ID No. 1) and pTaSU8A (Seq ID No. 6), a similarity of 96.8% results. Most of the differences regarding the sequences are in the 3'-untranslated region of the cDNAs. The remaining differences regarding the sequences in the coding region lead to different amine acids at a total of 12 positions of the derived protein sequences (Seq ID No. 3 and 7). The cDNAs contained in pTaSU19 and pTaSU8A are not identical and encode isoforms of the wheat isoamylase.

10

Example 6: Generation of the plant transformation vector pTa-alpha-SU8A

To express an antisense RNA corresponding to the TaSU8A cDNA, the plant transformation vector pTa-alpha-SU8A was constructed on the basis of the basic plasmid pUC19 by linking a part of the TaSU8A cDNA generated by PCR amplification in antisense orientation to the 3' end of the ubiquitin promoter. This promoter is composed of the first untranslated exon and the first intron of the maize *ubiquitin 1* gene (Christensen A.H. et al., Plant Mol. Biol 1 (1992), 675-689). Parts of the polylinker and the NOS terminator are derived from plasmid pAct1.cas (CAMBIA, TG 0063; Cambia, GPO Box 3200, Canberra ACT 2601, Australia). Vector constructs with this terminator and constructs based on pAct1.cas are described by McElroy et al. (Molecular Breeding 1 (1995), 27-37). The vector containing ubiquitin promoter, polylinker and NOS terminator and based on pUC19 was termed pUbi.cas.

To clone pTa-alpha-SU8A, an approx. 2.2 kb portion of the TaSU8A cDNA, viz. positions 140-2304 of Seq ID No. 6 was amplified by means of PCR.

30

The primers employed in this reaction were:

SUEX3: 5'-GCG GTA CCT CTA GAA GGA GAT ATA CAT ATG GCG GAG GAC AGG TAC GCG CTC-3' (Seq ID No. 10), and

SUEX4: 5'-GCT CGA GTC GAC TCA AAC ATC AGG GCG CAA TAC-3' (Seq ID No. 11).

1 ng of plasmid pTaSU8A was employed in the reaction as template. In addition, the PCR reaction contained: in each case 300 nM of the primers SUEX3 and SUEX4, in each case 200 µM of the nucleotides dATP, dGTP,

dCTP and dTTP, 1.6 mM MgCl₂, 60 mM Tris-SO₄ (pH 9.1), 18 mM (NH₄)₂SP₄ and 1 µl of Elongase[®] enzyme mix (mixture of Taq polymerase and DNA polymerase, Life Technologies). The final volume of the reaction was 50 µl. Amplification was performed in a PCR apparatus (TRIO[®]

- 5 Thermoblock, Biometra) with the following temperature regime: 1 min at 94°C (once); 30 sec at 95°C – 30 sec at 55°C – 2 min 30 sec at 68°C (30 cycles); 10 min at 68°C (once). The reaction gave rise to a DNA fragment 2205 bp in length.

- 10 The 2.2 kb product was restricted with *KpnI* and *Sall* and ligated into the expression vector pUbi.cas which had previously been cleaved with *KpnI* and *Sall*. The resulting plant transformation vector was termed pTa-alpha-SU8A and used as described above for transforming wheat.

We claim:

1. A nucleic acid molecule encoding a protein with the function of a wheat isoamylase, selected from the group consisting of
- 5 (a) a nucleic acid molecule encoding a protein comprising the amino acid sequence shown under Seq ID No. 3,
- (b) a nucleic acid molecule comprising the nucleotide sequence shown under Seq ID No. 2 or a part thereof or a ribonucleotide sequence corresponding hereto;
- 10 (c) a nucleic acid molecule which hybridizes with a nucleic acid molecule mentioned under (a) or (b) or is complementary thereto, and
- (d) a nucleic acid molecule whose nucleotide sequence deviates from the sequence of a nucleic acid molecule mentioned
- 15 under (a), (b) or (c) owing to the degeneracy of the genetic code,
- the nucleic acid molecule mentioned under (a), (c) and (d) having a homology of over 90% with Seq ID No. 2.
- 20 2. A nucleic acid molecule as claimed in claim 1, which is a DNA molecule.
3. A DNA molecule as claimed in claim 2, which is a cDNA molecule.
- 25 4. A nucleic acid molecule as claimed in one or more of claims 1 to 3 containing regulatory elements.
5. A nucleic acid molecule as claimed in claim 1, which is an RNA molecule.
- 30 6. A nucleic acid molecule which specifically hybridizes with a nucleic acid molecule as claimed in any of claims 1 to 5 and has a homology of over 90% with Seq ID No. 2.
- 35 7. A nucleic acid molecule as claimed in claim 6 which, is an oligonucleotide with a length of at least 15 nucleotides.

AMENDED SHEET

8. A vector containing a DNA molecule as claimed in any of claims 1 to 5.
- 5 9. A vector as claimed in claim 8, wherein said nucleic acid molecule is linked in sense orientation to regulatory elements which ensure transcription and synthesis of a translatable RNA in pro- or eukaryotic cells.
- 10 10. A vector as claimed in claim 8, wherein said nucleic acid molecule is linked in sense orientation to regulatory elements which ensure the synthesis of an untranslatable RNA in pro- or eukaryotic cells.
- 15 11. A vector as claimed in claim 8, wherein said nucleic acid molecule is linked in antisense orientation to regulatory elements which ensure the synthesis of an untranslatable RNA in pro- or eukaryotic cells.
- 20 12. A host cell which is transformed with a nucleic acid molecule as claimed in one or more of claims 1 to 5 or a vector as claimed in one or more of claims 8 to 11 or which is derived from such a cell.
- 25 14. A process for the preparation of a protein as claimed in claim 13, wherein a host cell as claimed in claim 12 is cultured under conditions which permit said protein to be synthesized and said protein is isolated from the cultured cells and/or the culture medium.
- 30 15. A process for generating a transgenic plant cell, wherein
a) a nucleic acid molecule as claimed in one or more of claims 1 to 5
or
b) a vector as claimed in one or more of claims 8 to 11
is integrated into the genome of a plant cell.
- 35 16. A transgenic plant cell which has been transformed with a nucleic acid molecule as claimed in one or more of claims 1 to 4 or one or more vector as claimed in claim 8 to 11 or which is derived from such a cell.
17. A process for generating a transgenic plant cell, wherein

a1) a nucleic acid molecule as claimed in one or more of claims 1 to 5 or

a2) a vector as claimed in one or more of claims 8 to 11

is integrated into the genome of a plant cell and

5 b) an intact plant is regenerated from said plant cell.

18. A plant containing a plant cell as claimed in claim 16.

19. A plant as claimed in claim 18, which is a crop plant.

10

20. A plant as claimed in claim 19, which is a starch-storing plant.

21. A plant as claimed in claim 20, which is a monocotyledonous plant or maize

15

22. A plant as claimed in claim 21, which is a barley, rye or wheat plant.

23. A propagation material of a plant as claimed in one or more of claims 18 to 22.

20

24. The use of a plant cell as claimed in claim 16, a plant as claimed in one or more of claims 18 to 22 or of propagation material as claimed in claim 23 for the production of starch.

95/M 206 WO

Abstract of the disclosure

Nucleic acid molecules encoding wheat enzymes involved in starch synthesis

Nucleic acid molecules are described which encode enzymes involved in starch synthesis in plants. These enzymes are wheat isoamylases.

The invention furthermore relates to vectors and host cells which contain the above-described nucleic acid molecules, in particular to transformed plant cells and plants which can be regenerated from these and which have an increased or reduced activity of the isoamylases according to the invention.

09674817 - 10600

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Nucleic acid molecules which code for enzymes derived from wheat and which are involved in the synthesis of starch

the specification of which

- is attached hereto
- was filed on May 07, 1999 as International Application PCT/EP99/03141 and including all the amendments through the date hereof.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s) for which Priority is Claimed:

Federal Republic of Germany, 19820608.9 of May 8, 1998

And I hereby appoint

William F. Lawrence, Registration No. 28,029, of the firm FROMMER LAWRENCE & HAUG, LLP whose post office address is 745 Fifth Avenue, New York, New York 10151, or their duly appointed associate, my attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to file continuation and divisional applications thereof, to receive the Patent, and to transact all business in the Patent and Trademark Office and in the Courts in connection therewith, and specify that all communications about the application are to be directed to the following correspondence address:

William F. Lawrence, Esq.
c/o FROMMER, LAWRENCE & HAUG LLP
745 Fifth Avenue
New York, New York 10151

Direct all telephone calls
to: (212) 588-0800, to the
attention of:
William F. Lawrence

09674937 110600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

INVENTOR(S) / Residence

1) Prof. Dr. Horst Loerz, Ramckeweg 6a, 22589 Hamburg, Germany

Signature: Mr. Horst Loerz

Date: 29.9.2000

2) Dr. Stephanie Luetticke, Lange Reihe 22, 20099 Hamburg, Germany

Signature: St. Luetticke

Date: 28.3.2000

3) Dr. Gernot Abel, Marstalsgade 6 4th, 2100 Kobenhaven, Denmark

Signature: _____

Date: _____

4) Dr. Ulrich Genschel, Eimsbütteler Str. 100b, 22 769 Hamburg, Germany

Signature: Ulrich Genschel

Date: 29.9.2000

1), 2) and 4) German

3) Austrian

Post Office Address of all Inventors:

Aventis CropScience GmbH
Patent- und Lizenzabteilung
Industriepark Höchst, Geb. K 801
D-65926 Frankfurt
Germany

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY

As below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

Nucleic acid molecules which code for enzymes derived from wheat and which are involved in the synthesis of starch

the specification of which

- is attached hereto
- was filed on May 07, 1999 as International Application PCT/EP99/03141 and including all the amendments through the date hereof.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s) for which Priority is Claimed:

Federal Republic of Germany, 19820608.9 of May 8, 1998

And I hereby appoint

William F. Lawrence, Registration No. 28,029, of the firm FROMMER LAWRENCE & HAUG, LLP whose post office address is 745 Fifth Avenue, New York, New York 10151, or their duly appointed associate, my attorneys, with full power of substitution and revocation, to prosecute this application, to make alterations and amendments therein, to file continuation and divisional applications thereof, to receive the Patent, and to transact all business in the Patent and Trademark Office and in the Courts in connection therewith, and specify that all communications about the application are to be directed to the following correspondence address:

**William F. Lawrence, Esq.
c/o FROMMER, LAWRENCE & HAUG LLP
745 Fifth Avenue
New York, New York 10151**

Direct all telephone calls
to: (212) 588-0800, to the
attention of:
William F. Lawrence

09674817 140600

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

INVENTOR(S) / Residence

1) Prof. Dr. Horst Loerz, Ramckeweg 6a, 22589 Hamburg, Germany

Signature: _____ Date: _____

2) Dr. Stephanie Luetticke, Lange Reihe 22, 20099 Hamburg, Germany

Signature: _____ Date: _____

3) ³⁻⁰⁰ Dr. Gernot Abel, Marstalsgade 6, 4th, 2100 Copenhagen, Denmark

Signature:  Date: 20/9-00
DEX

4) Dr. Ulrich Genschel, Eimsbütteler Str. 100b, 22 769 Hamburg, Germany

Signature: _____ Date: _____

1), 2) and 4) German

3) Austrian

Post Office Address of all Inventors:

Aventis CropScience GmbH
Patent- und Lizenzabteilung
Industriepark Höchst, Geb. K 801
D-65926 Frankfurt
Germany